



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

DEATHS DURING WEEK ENDED MARCH 29, 1919, IN CITIES.

The following table shows the registered deaths from all causes, and from pneumonia (all forms) and influenza combined, in certain large cities of the United States during the week ended March 29, 1919. The annual death rates per 1,000 population for the week and for the corresponding week of previous years are also shown.

The data are taken from the "Weekly Health Index," April 1, 1919, issued by the Bureau of the Census, Department of Commerce. The populations used in computing the rates are estimated by the Bureau of the Census as of July 1, 1918.

Registered deaths and annual death rates per 1,000 population in certain large cities of the United States, week ended Mar. 29, 1919—Deaths from all causes, and from pneumonia (all forms) and influenza combined.

| City. | Population July 1, 1918, estimated. | Total deaths, all causes. | Annual death rate per 1,000. | Annual death rate for preceding years. ¹ | Influenza and pneumonia (all forms). | |
|---------------------|--|------------------------------------|---------------------------------------|---|--|------------------------------------|
| | | | | | Number of deaths. | Annual death rate per 1,000. |
| Albany, N. Y. | 112,565 | 52 | 24.1 | C 20.4 | 12 | 5.6 |
| Atlanta, Ga. | 201,732 | 46 | 11.9 | C 16.0 | | |
| Baltimore, Md. | 669,981 | 212 | 16.5 | A 22.2 | 33 | 2.6 |
| Boston, Mass. | 785,245 | 239 | 15.9 | A 18.0 | 38 | 2.5 |
| Buffalo, N. Y. | 473,229 | 154 | 17.0 | C 19.8 | 22 | 2.4 |
| Cambridge, Mass. | 111,432 | 21 | 9.8 | A 16.2 | 3 | 1.4 |
| Chicago, Ill. | 2,596,681 | 733 | 14.7 | A 17.3 | 199 | 4.0 |
| Cincinnati, Ohio | 418,022 | 178 | 22.2 | C 19.5 | 59 | 7.4 |
| Cleveland, Ohio | 810,306 | 265 | 17.1 | C 13.9 | 98 | 6.3 |
| Columbus, Ohio | 225,296 | 82 | 19.0 | C 15.3 | 27 | 6.2 |
| Dayton, Ohio | 130,655 | 57 | 22.7 | C 16.4 | 14 | 5.6 |
| Denver, Colo. | | 81 | | | | |
| Fall River, Mass. | 128,892 | 36 | 14.6 | C 17.9 | 12 | 4.9 |
| Grand Rapids, Mich. | 135,450 | 30 | 11.6 | C 13.1 | | |
| Indianapolis, Ind. | 289,577 | 135 | 24.3 | C 18.7 | | |
| Jersey City, N. J. | 318,770 | 122 | 20.0 | C 22.7 | | |
| Kansas City, Mo. | 313,785 | 115 | 19.1 | C 27.3 | 44 | 7.3 |
| Los Angeles, Cal. | 568,495 | 160 | 14.7 | A 13.7 | 21 | 1.9 |
| Louisville, Ky. | 242,707 | 125 | 26.9 | C 20.6 | 45 | 9.7 |
| Lowell, Mass. | 109,681 | 37 | 17.7 | A 18.3 | 5 | 2.4 |
| Memphis, Tenn. | 154,759 | 78 | 26.3 | C 27.6 | 17 | 5.7 |
| Milwaukee, Wis. | 453,481 | 134 | 15.4 | A 16.1 | 41 | 4.7 |
| Minneapolis, Minn. | 553,442 | 112 | 15.2 | C 12.9 | | |
| Nashville, Tenn. | 119,215 | 43 | 18.8 | C 21.9 | 12 | 5.2 |
| Newark, N. J. | 428,684 | 155 | 18.9 | C 23.2 | 41 | 5.0 |
| New Haven, Conn. | 154,865 | 35 | 11.8 | C 18.9 | 7 | 2.4 |
| New Orleans, La. | 582,273 | 168 | 14.7 | A 21.9 | 20 | 2.7 |
| New York, N. Y. | 5,215,879 | 1,751 | 17.5 | C 23.8 | 516 | 5.2 |
| Oakland, Cal. | 214,266 | 50 | 12.2 | A 11.2 | | |
| Omaha, Nebr. | 180,264 | 43 | 12.5 | C 9.5 | | |
| Philadelphia, Pa. | 1,761,371 | 583 | 17.3 | A 18.6 | 127 | 3.8 |
| Pittsburgh, Pa. | 593,303 | 256 | 22.5 | C 22.2 | 108 | 9.5 |
| Portland, Oreg. | | 71 | | | 9 | |
| Providence, R. I. | 263,613 | 74 | 14.6 | C 27.7 | 15 | 3.0 |
| Richmond, Va. | 160,719 | 64 | 20.8 | C 16.5 | 11 | 3.6 |
| Rochester, N. Y. | 264,856 | 76 | 15.0 | C 15.0 | 14 | 2.8 |
| St. Louis, Mo. | 779,951 | 252 | 16.8 | C 17.4 | 77 | 5.1 |
| St. Paul, Minn. | 257,699 | 74 | 15.0 | C 16.8 | | |
| San Francisco, Cal. | 478,530 | 144 | 15.7 | C 20.4 | 15 | 1.6 |
| Seattle, Wash. | | 74 | | | 10 | |
| Spokane, Wash. | | 42 | | | | |
| Syracuse, N. Y. | 161,404 | 40 | 12.9 | C 19.7 | 5 | 1.6 |
| Toledo, Ohio | 262,234 | 101 | 20.1 | A 15.3 | 44 | 8.7 |
| Washington, D. C. | 401,681 | 114 | 14.8 | A 21.5 | 30 | 3.9 |
| Worcester, Mass. | 173,650 | 67 | 20.1 | C 19.5 | 12 | 3.6 |

¹ "A" indicates that the rate given is the average annual death rate per 1,000 population for the corresponding week of the years 1913 to 1917, inclusive. "C" indicates that the rate is the annual death rate per 1,000 population for the corresponding week of 1918.

² Population estimated as of July 1, 1919.

³ Rate is based on statistics of 1915, 1916, and 1917.